

Ferromagnetic ground states of the Hubbard model on a complete graph

Mario Salerno *

Department of Theoretical Physics
University of Salerno
84100 Salerno, Italy
and
Istituto Nazionale di Fisica della Materia

February 9, 2008

Abstract

We use group theory to derive the exact analytical expression of the ferromagnetic ground states of the Hubbard model on a complete graph for arbitrary lattice sites f and for arbitrary fillings N . We find that for $t > 0$ and for $N = f + 1$ the ground state is maximally ferromagnetic with total spin $S = (f - 1)/2$. For $N > f + 1$ the ground state is still ferromagnetic but becomes degenerate with respect to S .

*Email: SALERNO@csied.unisa.it

An interesting aspect of the theory of itinerant magnetism is the interplay between the kinetic energy and the potential energy of the electrons as a mechanism for the formation of ferromagnetic ground states. Examples of such mechanism leading to saturated ferromagnetism come from the Hubbard model with one hole in the strong coupling limit [1] and from the Hubbard model with finite coupling but on special graphs like the kagome lattice [2, 3]. The aim of the present paper is to show that ferromagnetic ground states exist also in the Hubbard model on a complete graph (i.e. with unconstrained hopping) for fillings higher than half filling and for arbitrary couplings. The unconstrained hopping implies the invariance of the Hamiltonian under the symmetric group S_f (f being the number of lattice sites), a feature which makes the model solvable for clusters of arbitrary sizes [4]. In spite of this simplification, the model keeps all the complexity of the standard Hubbard model coming from the Pauli exclusion principle and from the $SU(2)$ invariance. We hope that the knowledge of the exact ground states of this model may be useful for understanding the local properties of ferromagnetism in strongly correlated Fermi systems. In a previous paper we have shown that for fillings N below or equal to half filling ($N \leq f$), the ground states are always non ferromagnetic i.e. they are singlets or doublets depending on N being even or odd [4]. Here we extend this work by showing that just above half filling ($N = f + 1$) the ground state is not degenerate with respect to the total spin S and it is maximally ferromagnetic with $S = (f - 1)/2$. For higher fillings ($N > f + 1$), ferromagnetic ground states exist but they are degenerate with respect to S . This implies the existence of an hidden symmetry which resemble the well known spin-charge symmetry of the standard Hubbard model [6]. To analyze the system we extend the conjecture of ref. [4] on the symmetry properties of the ground states under the action of S_f and $SU(2)$, to the case of above half filling. This conjecture was derived from computer analytical diagonalizations of clusters of finite sizes by means of symbolic programs. Using this conjecture we derive analytical expressions for the ground states energies as a function of f and N and we show that the ferromagnetic ground states are always conducting. Our analytical results are found in excellent agreement with numerical diagonalizations this confirming the validity of our approach. The Hubbard model on a complete graph is written as

$$H = -t \sum_{\sigma, i \neq j}^f c_{i\sigma}^\dagger c_{j\sigma} + U \sum_i^f n_{i\uparrow} n_{i\downarrow}, \quad (1)$$

where $c_{i\sigma}^\dagger, c_{j\sigma}$, ($\sigma = \uparrow$ or $\downarrow$) are usual fermionic creation and annihilation operators. In ref. [4] we have shown how to use the representation theory of the permutation group to construct highest weight vectors of $SU(2)$ with definite S_f symmetry (for generalizations to arbitrary subgroups of S_f see ref.[5]). Here we just recall the main ideas leaving the details to ref.[4]. Let us denote with the symbols $|3\rangle, |2\rangle, |1\rangle, |0\rangle$ the four states on a given site (i.e. respectively, the doubly occupied, the single occupied spin up and spin down states and the vacuum) and let us introduce the quantum number

$$M = N + N_\uparrow. \quad (2)$$

To construct the basis functions which span the irreducible representations of the groups S_f and $SU(2)$ we consider all possible partitions $(m_1, m_2, \dots, m_f)$ of M into f parts (compatible with $N, N_\uparrow$), with $m_i = 0, 1, 2, 3$ and with $m_1 \geq m_2 \geq \dots \geq m_f$. Eigenmanifolds of S^2, S_z and S_f are then obtained by filling the quanta m_i of each partition in the boxes of the Young tableaux according to the following rules.

- i) The quanta must not increase when moving from left to right in a row or when moving down in a column.
- ii) The quanta referring to spin up and spin down states ($m_i = 1, 2$) must not appear more than once in a row.
- iii) The quanta referring to doubly occupied states or to empty states ($m_i = 3, 0$) must not appear more than once in a column.

In order to select among the filled tableaux the states corresponding to highest weight vectors of $SU(2)$ we use the following criterion. The application of the $SU(2)$ rising operator S^+ to a state changes a spin down into a spin up electron (1-2 flip). This implies that the states associated with tableaux satisfying the filling rules also after 1-2 flips are necessary states with $S > S_z$. The remaining tableaux will be automatically annihilated by S^+ i.e. they will give highest weight vectors of $SU(2)$. By implementing the above rules on a computer we have solved in exact form clusters of several sizes. From these studies the following conjecture is derived for the case $t > 0, U > 0$.

Conjecture: For $N = f + 1$ the ground state is maximally ferromagnetic with $S = (f - 1)/2$, it is non degenerate and it belongs to the completely antisymmetric representation $\{1^f\}$. For $N \geq f + 2$, the ground state is L -fold degenerate with respect to S where L is given by

$$L = \frac{2f - N}{2} + \delta, \quad (3)$$

with $\delta = 1$ or $\delta = \frac{1}{2}$ depending if N is even or odd. The spins S_j of the degenerate ground states are

$$S_j = \frac{2(f+1) - N}{2} - (j+1), \quad j = 0, 1, \dots, L-1, \quad (4)$$

and the corresponding S_f symmetries are given by tableaux T_j of type

$$T_j \equiv \{N-f, 2^j, 1^{2(f-j)-N}\}, \quad j = 0, 1, \dots, L-1. \quad (5)$$

Using this conjecture we can derive analytical expressions for the ground states above half filling. Let us first consider the case just above half filling, $N = f+1$, where saturated ferromagnetism occurs. According to our conjecture the ground state is a fully antisymmetric state with $S = (f-1)/2$. In order to be an highest weight vector of $SU(2)$ it must have $N_\uparrow = f$ and $N_\downarrow = 1$ so that from Eq. (2) we have $M = 2f+1$. There is only one way to fill $2f+1$ quanta into a tableaux of type $\{1^f\}$ according to our rules i.e. by putting a double occupied state 3 at the top and $f-1$ spin up states 2 in the remaining boxes as shown in Fig.1. By applying the antisymmetrizer operator to this tableau we obtain the maximally ferromagnetic ground state as

$$|\psi\rangle = \sum_{k=1}^f |k\rangle, \quad (6)$$

where $|k\rangle$ denotes a state with $f-1$ spin up and one double occupied state at site k . A direct calculation shows that $|\psi\rangle$ is an eigenstate of H with eigenvalue $-[t(N+2) - U]$

$$H|\psi\rangle = -(t(N+2) - U)|\psi\rangle. \quad (7)$$

A similar analysis can be performed for higher fillings. We have that for $N > f+1$ the condition for highest weight vectors is

$$2(f-j) - N = N_\uparrow - N_\downarrow, \quad j = 0, 1, \dots, L-1, \quad (8)$$

which together with $N = N_\uparrow + N_\downarrow$ gives the following expression for the quanta in Eq. (2) to be filled in tableaux T_j :

$$M_j = N + f - j, \quad j = 0, 1, \dots, L-1. \quad (9)$$

It is remarkable that each tableau of type T_j can be filled with M_j quanta just in one manner. We shall illustrate this with a concrete example. Let us

take $f = 5$ and $N = 7$. In this case we have $L = 2$ degenerate ground states $|\psi_i\rangle$, $i=0,1$, of spin $S_0 = 3/2$, $S_1 = 1/2$ and given by

$$\psi_0 = |33222\rangle - |23322\rangle - |23232\rangle - |23223\rangle, \quad (10)$$

$$\begin{aligned} \psi_1 = & |33221\rangle - |12233\rangle + |33122\rangle - |22133\rangle + \\ & |32231\rangle - |13223\rangle + |12332\rangle - |23321\rangle + \\ & |23123\rangle - |32132\rangle + |22331\rangle - |13322\rangle + \\ & 2(|33212\rangle - |21233\rangle + |23213\rangle - |31232\rangle + \\ & |21332\rangle - |23312\rangle). \end{aligned} \quad (11)$$

The corresponding filled tableaux are the ones reported in Fig. 2. The energy of these states is readily obtained as $E = \langle \psi_i | H | \psi_i \rangle$, this giving $E = -3t + 2U$. By extending this approach to arbitrary f, N we find that the energy of the ground state are given (for $N > f$ and $t > 0$) by

$$E(N, f) = -t(2f - N) + U(N - f). \quad (12)$$

We have checked Eq. (12) by comparing with Lanczos numerical diagonalizations of clusters of several sizes up to $f=16$, always obtaining a perfect agreement. The fact that for $N > f + 1$ the ferromagnetic ground states are degenerate with respect to S implies the existence of an hidden symmetry which we can describe as follows. The flipping of a single occupied state from spin up into spin down does not cost energy to the system if it is accompanied by a spatial rearrangement of the electrons i.e. if the electrons rearrange according to the next T_j symmetry given by Eq. (5). In Fig. 3 we report the filled tableaux T_j for fillings of type $N = f + k$ with k even (for k odd one proceeds similarly). We note that k is just the number of double occupied states and $0 \leq j \leq L - 1$ is the number of singly occupied spin down states present in the ground states. The tableaux T_0 always gives the maximally ferromagnetic ground state for a given filling. The others ground states are obtained from T_0 by removing spin up electrons from the first column and adding them as spin down electrons in the second column. Note that this rearrangement of the quanta is unique under the requirement of decreasing S while keeping the states highest weight vectors of $SU(2)$. From this we see that for $N = f + 1$ the ferromagnetic ground state cannot be degenerate (T_1 cannot be an highest weight vector). This accidental symmetry resemble the spin-charge symmetry observed on bipartite lattices. To clarify this let us recall that the spin-charge symmetry is related to the

partial particle-hole transformation

$$c_{i\uparrow} \rightarrow c_{i\uparrow} \quad (13)$$

$$c_{i\downarrow} \rightarrow (-1)^i c_{i\downarrow}^\dagger \quad (14)$$

which maps the $SU(2)$ algebra into the pseudo-spin $SU(2)$ algebra [6]. We note that the Hamiltonian (1) is not invariant under this transformation and the pseudo-spin does not commute with H . In spite of this, we have that the above ground states are highest weight vectors of the pseudo-spin algebra so that they are highest weight vectors of $SU(2) \times SU(2)$. To prove this statement we recall that the generators of the pseudo-spin algebra are

$$J^+ = \sum_i (-1)^i c_{i\uparrow}^\dagger c_{i\downarrow}^\dagger \quad (15)$$

$$J^- = \sum_i (-1)^i c_{i\downarrow} c_{i\uparrow} \quad (16)$$

$$J_z = \frac{N - f}{2}. \quad (17)$$

From the filled tableaux in Fig.3 we see that the ground states have no empty sites so that they are all annihilated by J^+ . Moreover, from

$$J^2 = J^- J^+ + J_z^2 + J_z \quad (18)$$

and from $N = f + k$ we see that they are eigenstates of J^2 with $J = J_z = k/2$. The fact that these ground states are invariant under pseudo-spin rotations whilst the Hamiltonian is not, shows how extending far the consequence of the Pauli exclusion principle could be. We also remark that by a standard particle-hole transformation the hamiltonian $H(t, U)$ goes into $H(-t, U) + (N - f)U$. This implies that the above ground states for $t, U > 0$ and $N > f$, can be obtained from the highest energy states of $f - N$ holes simply by changing the sign to t and shifting the energies by $(N - f)U$. Finally note that the ferromagnetic ground states are all conducting since they satisfy the equation

$$E(N - 1, f) - 2E(N, f) + E(N + 1, f) = 0, \quad (19)$$

as can be seen from Eq.12. This result is in agreement with the thermodynamical analysis performed in ref. [7].

Acknowledgments

Financial support from the Istituto Nazionale di Fisica della Materia, Unita' di Salerno, and from INTAS grant N. 93-1324 is acknowledged.

Figure Captions

Fig.1 Filled Young tableaux corresponding to the ferromagnetic ground state of spin $S = \frac{f-1}{2}$ at filling $N = f + 1$.

Fig.2 Filled Young tableaux for the degenerate ground states of the case $f = 5$, $N = 7$. The total spin S of these states is respectively (from left to right) $3/2$, $1/2$.

Fig.3 Filled Young tableaux of type T_j associated with degenerate ground states at filling $N = f + k$ with k even, $S_j = (f - 2j - k)/2$ and pseudo-spin $J = k/2$. Here k denotes the number of double occupied states and j the number of singly occupied spin down states present in the ground states.

References

- [1] Y Nagaoka, *Phys.Rev.B*,147 (1966) 392.
- [2] A Mielke, *J.Phys.A*,25 (1992) 4335.
- [3] H Tasaki, *Phys.Rev.Lett.*, 69 (1992) 1608.
- [4] M Salerno, *Z.Phys.B*, (1996) in press.
- [5] M Salerno, *Phys.Lett.A*, (1996) in press.
- [6] F Woynarovich, *J.Phys.C*, 16 (1983) 6593.
- [7] P van Dongen, D Volhardt. *Phys.Rev.B*, 40 (1989) 7252.

3
2
$\vdots$
2

Fig.1




Fig.2




Fig.3